

Original Article

**ASSESSMENT OF PATIENT COUNSELLING IN DIABETIC AND HYPERTENSIVE PATIENTS IN TERMS OF PATIENT KNOWLEDGE ABOUT THEIR MEDICATION, DISEASE STATE AND LIFESTYLE MODIFICATIONS BY PHARMACY PERSONNEL AT LDF CLINIC IN MASERU**

LINEO MAJA<sup>1</sup>, THABISO MASIA<sup>2</sup>, KABELO BINYANE<sup>2</sup>, MASEABATA RAMATHEBANE<sup>3</sup>

<sup>1</sup>Lecturer, Department of Pharmacy, National University of Lesotho, <sup>2</sup>Student, Department of Pharmacy, National University of Lesotho,

<sup>3</sup>Senior Lecturer, Department of Pharmacy, National University of Lesotho

Email: majalineo@gmail.com

Received: 15 Dec 2017 Revised and Accepted: 13 Feb 2018

**ABSTRACT**

**Objective:** To investigate if diabetic and hypertensive patients were being adequately counselled by pharmacy personnel about their medication, disease states and lifestyle modifications at Lesotho defence force (LDF) clinic in Maseru.

**Methods:** A cross-sectional study was conducted among diabetic and hypertensive patients treated at LDF clinic in Maseru from March to May 2017. A semi-structured questionnaire was used to collect data in a face-to-face interview with patients. Microsoft Excel 2010 was used to analyse collected data.

**Results:** The study population consisted of 222 participants; 22 (28.2%) of diabetic respondents had inadequate knowledge of diabetes mellitus complications and 56 (71.8%) had no knowledge of complications. 153 (78.8%) of hypertensive respondents had inadequate knowledge of hypertension complications and 33 (17.1%) had no knowledge of complications. 56 (25.2%) had adequate knowledge of lifestyle modifications of diabetes mellitus and hypertension and 164 (73.9%) had inadequate knowledge. 182 (82.0%) had adequate knowledge of their anti-hypertensive and anti-diabetic medicines names, 20 (9.0%) had inadequate knowledge and 20 (9.0%) had no knowledge. 63 (28.4%) had adequate knowledge of their medicines strengths, 17 (7.7%) had inadequate knowledge and 142 (64.0%) had no knowledge. 199 (89.6%) respondents had adequate knowledge of dosing frequency and 20 (9.0%) had inadequate knowledge.

**Conclusion:** Hypertensive and diabetic patients' knowledge about medication, disease states and lifestyle modifications were inadequate due to poor patient counselling on such aspects by pharmacy personnel at LDF clinic in Maseru.

**Keywords:** Diabetes mellitus, Hypertension, Patient counselling, Medication knowledge, lifestyle modifications, Pharmacy personnel

© 2018 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) DOI: <http://dx.doi.org/10.22159/ijpps.2018v10i3.24291>

**INTRODUCTION**

Non-communicable diseases (NCDs) contributed to the rise in mortality in the world. According to the World Health Organization (WHO) [1], 38 million people died annually of NCDs with cardiovascular diseases responsible for 17.5 million deaths, cancer 8.2 million, respiratory diseases 4 million, and diabetes 1.5 million deaths. The 2014 World Health Organization NCDs national profile of Lesotho reported that 12% of deaths were due to cardiovascular diseases, 3% diabetes mellitus and 3% due to respiratory diseases [2]. The World Health Organization [3] stated that 27% of deaths in Lesotho were due to NCDs.

Patient knowledge on diabetes mellitus management includes medicine use, risk factors associated with diabetes mellitus, and lifestyle modifications [4] which are important factors in pharmaceutical care of diabetic patients. Adequate patient knowledge on their diseases, complications associated with those diseases, and drug therapy can improve self-management which forms a vital part of an effective treatment plan [4] thus, improving patient compliance to treatment and decreasing complications associated with the disease [5]. Furthermore, patient knowledge of blood pressure (BP) in hypertension, blood glucose control in diabetes mellitus, and medications adherence play an important role in the management of diabetes mellitus and hypertension [6, 7].

Patient counselling is an effective method to improve patient medication adherence [7]. Counselling involves patient education on diseases and their complications, medication use and storage, and lifestyle modifications [7, 8]. Pharmacy personnel have to encourage patient understanding on how the prescribed treatment works with emphasis on medicines dosing frequency, route of administration

and side effects [6]. Benefits of patient counselling include improved patient understanding of their diseases and its treatment, adherence, and improved therapeutic outcome [6, 7]. Additionally, pharmacists are able to gain knowledge of medication use from the patient's perspective [7]. Educating patients about their disease states and medication can also increase their active participation in therapy and improve medication adherence [9].

Pharmacists are the third largest health profession in the world after doctors and nurses [10] and help to identify, assess, educate, refer and continuously monitor patients in diabetes mellitus and hypertension management. Pharmacists are in a unique position to play a vital role in helping patients to cope with their disease and make informed decisions regarding management and medication through patient education [4, 10]. Therefore, responsibilities of pharmacists involve long-term patient supervision, patient education activities, consideration of medication-related issues such as side effects, and optimisation of medicinal treatment and adherence [11].

The challenge to adequate patient counselling for hypertensive and diabetic patients is the increasing number of patients. Thus, the rationale was that hypertensive and diabetic patients counselling is essential to enable patients to care for themselves at home, handle medication-related problems, and to improve adherence so due to patient increase, insufficient time was dedicated to delivering patient counselling. Therefore, the objective of this study was to determine if hypertensive and/or diabetic patients were counselled on disease states, medication (names, strengths, dose frequency, side effects and storage), and lifestyle modifications by pharmacy personnel at LDF clinic in Maseru

**MATERIALS AND METHODS****Study setting**

The study was conducted at LDF clinic in Maseru district in Lesotho.

**Study design**

This was a cross-sectional study where data on hypertensive and/or diabetic patients knowledge regarding medication, disease states and lifestyle modification were collected from the study population at a single point in time (from March to May 2017) at LDF clinic in Maseru district in Lesotho.

**Target population**

The target population consisted of all male and female adult outpatients 18 y of age and above with either diabetes mellitus, hypertension, or with both hypertension and diabetes mellitus at LDF clinic.

**Sampling**

The preliminary number of diabetic and/or hypertensive patients at LDF clinic was 523 in 2017. The sample size was calculated at 95% confidence interval, with 5% margin of error. The calculated sample size used as the study population was 222 patients with diabetes mellitus and/or hypertension. No sampling method was used. Inclusion criteria were as follows: all diabetic and/or hypertensive outpatients (males and females) above 18 y of age at LDF clinic that have been on diabetes mellitus and hypertension treatment for more than a year. An exclusion criterion was hypertensive and/or diabetic patients who did not visit the LDF clinic during the days when the researchers were gathering data.

**Data collection**

A semi-structured questionnaire with both open and closed-ended questions was used as a data collection tool. An extensive literature review of similar studies was conducted during the development of the semi-structured questionnaire that assisted in coming up with unambiguous questions. Additionally, the following studies were used to formulate the semi-structured questionnaire: (a) Study of knowledge, attitude, and practice of general population of Guntur towards silent killer diseases: hypertension and diabetes by Bollu *et al.* [12], and (b) Influence of post-discharge counselling on health outcomes in diabetic and hypertensive patients by Adepu *et al.* [6].

**RESULTS****Table 1: Age, gender and disease state(s) distribution of respondent**

Age group (y)	Frequency (n)	Percentage (%)
21-30	16	7.2%
31-40	44	19.8%
41-50	63	28.4%
51-60	55	24.8%
61-70	35	15.8%
71-80	9	4.1%
<b>Total respondents</b>	<b>222</b>	<b>100.0%</b>
<b>Gender</b>	<b>Frequency (n)</b>	<b>Percentage (%)</b>
Male	46	20.7%
Female	176	79.3%
<b>Total respondents</b>	<b>222</b>	<b>100.0%</b>
<b>Disease state(s)</b>	<b>Frequency (n)</b>	<b>Percentage (%)</b>
Hypertension	144	64.9%
Diabetes mellitus	29	13.1%
Hypertension and diabetes mellitus	49	22.1%
<b>Total respondents</b>	<b>222</b>	<b>100.0%</b>

Table 1 shows that the majority (63, 28.4%) of respondents were in the age group range 41-50 and the majority (176, 79.3%) of respondents were female. Table 1 also indicates that about two thirds (144, 64.9%) of respondents had hypertension.

The questionnaire was prepared in both English and Sesotho languages. Hypertensive and/or diabetic patients attending the clinic were provided with information about the study and informed consent forms using their home language (Sesotho). Patients who gave their informed consent participated in the study. The researchers then conveniently conducted face-to-face interviews using the semi-structured questionnaire and looked through medication booklets of diabetic and/or hypertensive patients who attended the clinic during the days the researchers were gathering data at the clinic. The researchers checked the patients' medication booklets to confirm their diabetic and hypertensive states, anti-diabetic and anti-hypertensive medication, and to confirm that they met the inclusion criteria.

**Time period**

Data were collected from March to May 2017 using semi-structured questionnaires in a face-to-face interview. Data was collected on Tuesdays, Wednesdays and Thursdays every week between 08:00 and 16:00.

**Data analysis**

Data were analysed using Microsoft Excel 2010. Descriptive statistics (frequency, percentage, mean, and standard deviation) was used to demonstrate characteristics of the study population. The responses from participants regarding the complications of hypertension and diabetes mellitus, lifestyle modifications, and medicines (names, strengths and dosing frequency) were classified using knowledge scores. Participants were said to have "adequate" knowledge if they mentioned at least five correct answers; "inadequate" knowledge if they mentioned at most four correct answers and "no" knowledge (Don't know) if they did not know.

Furthermore, participants were said to have "adequate" knowledge if they responded with a percentage score of "75-100%"; "inadequate" knowledge "<75% but>0%" and "no" knowledge (Don't know) if they did not know anything, with a percentage score of "(0%)".

**Ethical approval and consent to participate**

Ethical approval was granted by the Local Research Ethics Committee of the National University of Lesotho (NUL) and the Ministry of Health Research and Ethics Committee of Lesotho, registration number 76-2017. Goodwill permission to conduct the study was obtained from the LDF clinic manager. The study participants gave their informed consent.

**Table 2: Distribution of respondents by highest level of education**

Highest level of education	Frequency (n)	Percentage (%)
Primary school	98	44.1%
Secondary school	70	31.5%
High school	29	13.1%
College	16	7.2%
University	7	3.2%
No education	2	0.9%
Total respondents	222	100.0%

According to table 2, the majority (98, 44.1%) of respondents had primary school level of education.

**Table 3: Age, fasting blood glucose (FBG) level and blood pressure (BP) means**

Variable	Mean ( $\pm$ sd)	Minimum	Maximum
Age (Years)	48.7 $\pm$ 12.0	21	77
Fasting Blood Glucose (mmol/l)	6.9 $\pm$ 2.0	3.9	14.5
Blood Pressure (mmHg)	Systolic BP      Diastolic BP	Systolic BP      Diastolic BP	Systolic BP      Diastolic BP
SD: Standard deviation	143.4 $\pm$ 13.8      89.6 $\pm$ 8.2	110                  51	194                  110

Table 3 indicates that the mean age of respondents was 48.7 $\pm$ 12.0 y, mean fasting blood glucose (FBG) level of diabetic respondents was 6.9 $\pm$ 2.0 mmol/l, mean systolic blood pressure (BP) of hypertensive patients was 143.4 $\pm$ 13.8 mmHg, and the mean diastolic BP was 89.6 $\pm$ 8.2 mmHg.

**Table 4: Knowledge regarding hypertension and diabetes mellitus**

Disease incurable	Frequency (n)	Percentage (%)
Yes	177	79.7%
No	0	0.0%
Don't know	45	20.3%
Total respondents	222	100.0%
Complications of diabetes mellitus	N=78	
	Frequency (n)	Percentage (%)
Retinopathy	9	11.5%
Diabetic foot	18	23.1%
Nephropathy	1	1.3%
Neuropathy	0	0.0%
Complications of hypertension	N=193	
	Frequency (n)	Percentage (%)
Stroke	160	82.9%
Kidney failure	8	4.1%
Heart attack	6	3.1%
Heart failure	10	5.2%

Table 4 shows that about two thirds (177, 79.7%) of respondents knew that the diseases were incurable. The majority (18, 23.1%) of respondents knew about a diabetic foot as a complication of diabetes mellitus. Additionally, the majority (160, 82.9%) of respondents knew that stroke was a complication for hypertension.

**Table 5: Knowledge regarding target blood pressure (BP) and fasting blood sugar (FBG) level**

Blood pressure (BP)	Frequency (n)	Percentage (%)
<140/90 mmHg	97	50.3%
$\geq$ 140/90 mmHg	0	0.0%
Don't know	96	49.7%
Total respondents	193	100.0%
Fasting blood sugar (FBG)	Frequency (n)	Percentage (%)
<6.2 mmol/l	28	35.9%
$\geq$ 6.2 mmol/l	0	0.0%
Don't know	50	64.1%
Total respondents	78	100.0%

Table 5 indicates that half (97, 50.3%) of respondents knew the target BP, and more than half (50, 64.1%) of respondents did not know the target FBG level.

**Table 6: Knowledge regarding lifestyle modifications of diabetes mellitus and hypertension**

Lifestyle modifications in diabetes mellitus and hypertension	N=222	
	Frequency (n)	Percentage (%)
Weight loss if obese	20	9.0%
Reduce intake of fatty foods	210	94.6%
Reduce alcohol consumption	48	21.6%
Smoking cessation	17	7.7%
Restrict salt intake	186	83.8%
Physical exercise	67	30.2%
Keeping stress under control	60	27.0%
Intake of fruits and vegetables	88	39.6%
Reduce sugar intake	37	16.7%

Table 6 shows that the majority (186, 83.8%) of respondents knew about salt intake restriction as one of the lifestyle modifications in diabetes mellitus and hypertension.

Table 7: Knowledge regarding hypertension and diabetes mellitus medication names and strengths

Hypertension medication names	Frequency (n)	Percentage (%)
Hydrochlorothiazide (N=192)	168	87.5%
Atenolol (N=109)	86	78.9%
Captopril (N=147)	118	80.3%
Hydralazine (N=2)	1	50.0%
Aspirin (N=103)	89	86.4%
Nifedipine (N=56)	37	66.1%
<b>Diabetes mellitus medication names</b>	<b>Frequency (n)</b>	<b>Percentage (%)</b>
Glibenclamide (N=29)	22	75.9%
Metformin (N=57)	53	93.0%
Insulin (N=49)	46	95.8%
<b>Hypertension medication strengths</b>	<b>Frequency (n)</b>	<b>Percentage (%)</b>
Hydrochlorothiazide 25 mg (N=192)	76	39.6%
Atenolol 50 mg and 100 mg (N=109)	44	40.4%
Captopril 25 mg and 50 mg (N=147)	56	38.1%
Hydralazine 25 mg (N=2)	0	0.0%
Aspirin 150 mg (N=103)	34	33.0%
Nifedipine 10 mg (N=56)	20	35.7%
<b>Diabetes mellitus medication strengths</b>	<b>Frequency (n)</b>	<b>Percentage (%)</b>
Glibenclamide 5 mg (N=29)	8	27.6%
Metformin 500 mg and 850 mg (N=57)	22	38.6%
Insulin 100 IU/ml (N=49)	1	2.1%

N=Total number of patients on that particular drug; n=patients who know the name of that particular drug.

Table 7 shows that the majority (168, 87.5%) of respondents on an anti-hypertensive drug, hydrochlorothiazide, knew its name, and the majority (46, 95.8%) of respondents on an anti-diabetic drug, insulin, knew its name. Table 7 also shows that the majority (44, 40.4%) of respondents on an anti-hypertensive drug, atenolol, knew its strength, and the majority (22, 38.6%) of respondents on an anti-diabetic drug, metformin, knew its strength.

Table 8 shows that the majority (187, 97.4%) of respondents on an anti-hypertensive drug, hydrochlorothiazide, knew its dosing frequency, and the majority (49, 100.0%) of respondents on an anti-diabetic drug, insulin, knew its dosing frequency. Additionally, table 8 shows that the majority (91, 41.0%) of respondents store their diabetes mellitus and/or hypertension medicines at room temperature.

Table 8: Knowledge regarding hypertension and diabetes mellitus medication dosing frequency and storage

Hypertension medication dosing frequency	Frequency (n)	Percentage (%)
Hydrochlorothiazide o. d. (N=192)	187	97.4%
Atenolol o. d. (N=109)	100	91.7%
Captopril b. d. and t. d. s. (N=147)	127	86.4%
Hydralazine o. d. (N=2)	1	50.0%
Aspirin o. d. (N=103)	96	93.2%
Nifedipine b. d. and t. d. s. (N=56)	47	83.9%
<b>Diabetes mellitus medication dosing frequency</b>	<b>Frequency (n)</b>	<b>Percentage (%)</b>
Glibenclamide o. d. and b. d. (N=29)	26	89.7%
Metformin b. d. and t. d. s. (N=57)	53	93.0%
Insulin b. d. (N=49)	49	100.0%
<b>Medication storage</b>	<b>Frequency (n)</b>	<b>Percentage (%)</b>
Room temperature (15 to 25° C)	91	41.0%
Cool temperature (8 to 15° C)	26	11.7%
Refrigeration (2 to 8° C)	18	8.1%
Freezing temperatures (-10 to -25° C)	0	0.0%
Warm temperatures (>25° C)	87	39.2%
<b>Total respondents</b>	<b>222</b>	<b>100.0%</b>

*o. d.-once daily, b. d.-twice daily, t. d. s.-three times daily*

N=Total number of patients on that particular drug; n=patients who know the dosing frequency of that particular drug.

Table 9: Respondents who experienced diabetes mellitus and/or hypertension medicines side effects and action taken

Medicine side-effects	Frequency (n)	Percentage (%)
Yes	116	52.3%
No	106	47.7%
<b>Total respondents</b>	<b>222</b>	<b>100.0%</b>
<b>Side effects experienced</b>	<b>Frequency (n)</b>	<b>Percentage (%)</b>
Dizziness	82	70.7%
Headache	19	16.4%
Fatigue	36	31.0%
Redness at injection side	5	4.3%
Cough	3	2.6%

Medical action taken (if side-effects experienced)	Frequency (n)	Percentage (%)
Consult doctor immediately	4	3.4%
Consult doctor on the next visit	8	6.9%
No action taken	102	87.9%
Stop taking the medicine	2	1.7%
<b>Total respondents</b>	<b>116</b>	<b>100.0%</b>
Hypoglycaemia symptoms management	Frequency (n)	Percentage (%)
Eat carbohydrates meal immediately	4	5.1%
Intake of sugar/sweets	60	76.9%
Eat carbohydrates meal immediately or intake of sugar/sweets	9	11.5%
Don't know	5	6.4%
<b>Total respondents</b>	<b>78</b>	<b>100.0%</b>

Table 9 shows that the majority (116, 52.3%) of respondents experienced diabetes mellitus and/or hypertension medicines side effects, 82 (70.7%) of respondents experienced dizziness as a side effect, and 102 (87.9%) of respondents who experienced medicines side effects did not take any medical action. Table 9 also shows that the majority (60, 76.9%) of respondents take sugar/sweets to manage hypoglycaemia symptoms.

**Table 10: Knowledge scores regarding medication, disease states and lifestyle modifications**

Knowledge scores	Frequency (n) and percentage (%)	Adequate ( $\geq$ 75%)	Inadequate (<75% but >0%)	Don't know (0%)	Total respondents
Complications of diabetes mellitus	n %	0 0.0%	22 28.2%	56 71.8%	78 100.0%
Complications of hypertension	n %	8 4.1%	152 78.8%	33 17.1%	193 100.0%
Lifestyle modifications	n %	56 25.2%	164 73.9%	2 0.9%	222 100.0%
Medication names	n %	182 82.0%	20 9.0%	20 9.0%	222 100.0%
Medication strengths	n %	63 28.4%	17 7.7%	142 64.0%	222 100.0%
Medication dosing frequency	n %	199 89.6%	20 9.0%	3 1.4%	222 100%

According to table 10, more than half (56, 71.8%) of respondents did not know any diabetes mellitus complications, and more than half (152, 78.8%) of respondents had inadequate knowledge of hypertension complications. About two thirds (164, 73.9%) of respondents had inadequate knowledge regarding lifestyle modifications in diabetes mellitus and hypertension. The majority (183, 82.0%) of respondents had adequate knowledge regarding their diabetes mellitus and/or hypertension medicines names, 142 (64.0%) of respondents did not know any of their diabetes mellitus and/or hypertension medicines strengths, and 199 (89.6%) of respondents had adequate knowledge regarding their diabetes mellitus and/or hypertension medicines dosing frequency.

## DISCUSSION

Our study aimed to assess if diabetic and/or hypertensive patients were being adequately counselled by pharmacy personnel about their medications, disease states and lifestyle modifications at LDF clinic in Maseru. The aim was achieved through determination of knowledge of such aspects among diabetic and hypertensive patients visiting the clinic. In our study, the majority of respondents were hypertensive and the majority of respondents were female. Most respondents were within the age range of 41-50 y of age. The mean age of respondents was 48.7±12.0 y. additionally, most respondents had primary school educational level.

Our study showed that the mean systolic BP and the mean diastolic BP for hypertensive patients were high. Therefore, hypertensive patients had high uncontrolled BP despite the fact that they were on anti-hypertensive medicines. In a study conducted by Akoko *et al.* in Cameroon, BP control was poor, with less than half of the participants (42.1%) having good BP [13] and these findings were consistent with the findings in our study. The normal systolic BP should be <120 mmHg and the normal diastolic BP should be <80 mmHg [14]. Also, our study showed that the mean FBG level was high for diabetic patients meaning glycaemic control was poor among diabetic patients even though they were taking anti-diabetic medicines. Our findings were consistent with those in a study by Dekker *et al.* which showed that inadequate glycaemic control was present in 74% of all diabetic patients [15]. Additionally, a study by Nepal *et al.* conducted in Nepal revealed that among 89 type 2 diabetic respondents only 27 (30.34%) had normal FBG level [16]. The Global Diabetes Community [17] stated that normal FBG level should be within a range of 4-7 mmol/l for people with type 1 or type 2 diabetes mellitus. The uncontrolled BP and blood glucose

levels, although the patients were on drug treatment, suggested that inadequate counselling on medicine use and lifestyle modification in controlling BP and the glycaemic level was not provided to patients by pharmacy personnel. A study conducted by Adepu *et al.* on post-discharge counselling in diabetic and hypertensive patients showed that there was improvement in BP and capillary blood glucose levels in test group patients compared to control group patients due to patient education on medication adherence behaviour, dietary changes adoption and lifestyle modifications [6].

In our study, half of the respondents knew the target BP and more than half of respondents did not know the target FBG level. Thus, pharmacy personnel at LDF clinic provided adequate patient counselling on target BP as opposed to target FBG level of which knowledge of target BP and FBG level is essential in self-monitoring of hypertension and diabetes mellitus, respectively. A study by Nepal *et al.* conducted in Nepal showed that 74.16% of patients were aware of the importance of regular checking of BP hence, knowledge of patients regarding regular self-monitoring of blood glucose level and BP is essential [16]. Patient knowledge and awareness of target BP and FBG levels play important roles in the ability to successfully self-monitor hypertension and diabetes mellitus.

Our study also revealed that about two-thirds of respondents knew that hypertension and diabetes mellitus were incurable. The findings of our study were consistent with those in a study by Almas *et al.* where 166 (76%) men and 162 (73.3%) women agreed that hypertension was a lifelong disease (p-value 0.3) [18]. Respondents in our study knew that hypertension and diabetes mellitus were incurable thus patient counselling provided by pharmacy personnel at LDF clinic was adequate with regards to hypertension and diabetes mellitus being curable or incurable. The findings in our

study were in contrast with those in a study by Familoni *et al.* indicating that one-third (35.4%) of the patients knew that hypertension should be treated for life [19]. The findings in a study by Fairchild *et al.* were also in contrast with our findings where about two-thirds (39, 65%) of participants believed that their diabetes could be cured [20].

Hypertensive and diabetic patients develop complications due to poor awareness regarding the diseases and inadequate BP and blood glucose level control. In our study, the majority of respondents knew about a diabetic foot as a complication of diabetes mellitus and stroke as a complication of hypertension. A study by Obirikorang *et al.* conducted in Ghana showed that type 2 diabetes mellitus complication commonly known by diabetic patients was a diabetic foot (51.5 %) [21], which was consistent with our study findings. Knowledge scores in our study showed that the majority of respondents did not know about diabetes mellitus complications and the majority had inadequate knowledge about hypertension complications. Our study findings were consistent with a study by Obirikorang *et al.* conducted in Ghana indicating that a higher proportion of type 2 diabetes patients did not have adequate knowledge on diabetic complications [21]. A study by Adepu *et al.* showed that at baseline the majority of participants were unaware of asymptomatic hypertension and complications associated with it [22]. Additionally, our study findings were consistent with a study by Abd El-Hay *et al.* where most patients (67.3%) had fair level of knowledge regarding symptoms and complications of hypertension [23]. Hypertensive and diabetic patients in our study were not aware of other complications of hypertension (kidney failure, heart attack, heart failure) and diabetes mellitus (retinopathy, nephropathy, neuropathy) suggesting that patients were not well informed about the complications of hypertension and diabetes mellitus during patient counselling by pharmacy personnel at the LDF clinic.

Our study also showed that most hypertensive and diabetic patients knew about salt intake restriction as one of the lifestyle modifications. A study by Abd El-Hay *et al.* on knowledge and perception related to hypertension and lifestyle behaviour modifications showed that the majority of patients knew about excessive salt intake increase (82.2%) as a risk for developing hypertension [23]; their findings were consistent with the findings in our study. The findings in a study by Bollampally *et al.* showed that 66% of participants thought that taking less salt will control the BP; their findings were similar to the findings in our study [24].

Additionally, knowledge scores in our study revealed that majority of hypertensive and diabetic patients had inadequate knowledge of lifestyle modifications. Our study findings were in contrast with findings in a study by Abd El-Hay *et al.* where most patients (67.3%) had a fair level of knowledge regarding lifestyle modifications to control hypertension [23]. In our study, hypertensive and diabetic patients had inadequate knowledge of lifestyle modifications (such as weight loss, fatty foods, alcohol and sugar intake reduction, physical exercise, stress management, and fruits and vegetables intake) in hypertension and diabetes mellitus management due to poor patient counselling on lifestyle modifications by pharmacy personnel at LDF clinic.

Patient counselling includes providing verbal and/or written medication information on directions of use, advice on side effects, and storage to patients. Our study revealed that hypertensive patients mostly knew hydrochlorothiazide by name and its dosing frequency while diabetic patients mostly knew insulin by name and its dosing frequency. The findings in our study also showed that the majority of hypertensive patients on atenolol knew its strength and the majority of diabetic patients on metformin knew its strength. Knowledge scores on anti-hypertensive and anti-diabetic medicines in our study showed that most respondents had adequate knowledge of their medicine names and dosing frequencies, and about two-thirds of respondents did not know their medicines strengths. Our findings were consistent with a study by Bollampally *et al.* indicating that patients had good knowledge of prescribed drug names [24]. A study by Nnaemeka *et al.* revealed that most diabetic patients (39, 54.2%) knew the names of their anti-diabetic

medications and the majority (52, 72.2%) of participants knew the dosing frequency of their anti-diabetic medicines [8]; their findings were similar to finding in our study. Adequate knowledge of medicine names and dosing frequencies as revealed by our study findings suggested that good counselling was provided by pharmacy personnel at LDF clinic on such aspects. On the other hand, there was poor knowledge regarding anti-hypertensive and anti-diabetic medicines strengths suggesting lack of patient counselling regarding medicines strengths.

The majority of respondents in our study experienced anti-hypertensive and anti-diabetic medicines side effects and the side effect mostly experienced by respondents was dizziness which could be due to uncontrolled levels of BP and FBG. Our findings were in contrast with those in a study by Manobharathi *et al.* where 90 (83.3%) diabetic patients on anti-diabetic medicines had no side effects with the medicine [25]. One of the factors affecting uncontrolled levels of BP and FBG in hypertensive and diabetic patients, respectively include non-compliance to medicines due to side effects. A study by Manobharathi *et al.* showed that experiencing side effects and reporting the side effects experienced were found to be more among non-compliant patients compared to compliant patients [25]. A study by Akoko *et al.* also revealed that participants who reported side effects of their drugs were significantly less compliant than those who had never experienced any side effects [13]. Additionally, our findings showed that most hypertensive and diabetic patients who experienced medicines side effects did not take any medical action. Our study findings suggested the serious need for pharmacy personnel to improve patient counselling on medication side effects as well as actions patients should take when experiencing these side effects. Also, most respondents in our study took sugar/sweets to manage hypoglycaemic symptoms thus hypertensive and diabetic patients knew how to manage hypoglycaemia. Our findings were comparable with findings from a study by Adepu *et al.* indicating that most of the patients were aware of how to recognise and control the hypoglycaemic symptoms [22]. Therefore, there was adequate counselling regarding the management of hypoglycaemic symptoms by pharmacy personnel at LDF clinic.

### Limitations

Some of the respondents were not eager to participate in the study because they thought their participation would affect the type of clinical services they were offered at the LDF clinic.

### CONCLUSION

There is an increasing burden of non-communicable diseases such as diabetes mellitus and hypertension in the world which requires patients to take medicines. Therefore, the interaction between diabetic and/or hypertensive patients and pharmacy personnel through patient counselling has an impact on adherence to therapy. Poor relationship between patients and pharmacy personnel results in poor adherence to medication which is associated with higher glucose levels for the diabetic patient and high blood pressure for hypertensive patients. In general, there is inadequate patient knowledge regarding diabetes mellitus and hypertension complications, target blood pressure and fasting blood glucose levels, lifestyle modifications, medicines strengths, and medical actions to take when experiencing side effects in diabetes mellitus and hypertension management due to poor patient counselling by pharmacy personnel at LDF clinic in Maseru, Lesotho. Our study has therefore revealed that hypertensive and diabetic patients lack valuable information about their medicines, disease states and lifestyle modification thus counselling strategies to strengthen patient counselling should be developed.

### AUTHORS CONTRIBUTIONS

Lineo Maja developed the study, collected and analysed the data, and wrote the first draft of the paper. Additionally, Lineo Maja read and approved the final manuscript.

Thabiso Masia developed the study, collected and analysed the data, and wrote the first draft of the paper.

Kabelo Binyane developed the study, collected and analysed the data, and wrote the first draft of the paper.

Maseabata Ramathebane read and approved the final manuscript.

#### CONFLICTS INTERESTS

All authors declare that they have no conflict of interest

#### REFERENCES

- World Health Organization. Non-communicable diseases fact sheet; 2015. <http://www.who.int/mediacentre/factsheets/fs355/en/>. [Last accessed on 11 Apr 2016].
- World Health Organization. Non-communicable diseases (NCDs) country profile; 2014. [http://www.who.int/nmh/countries/lso\\_en.pdf](http://www.who.int/nmh/countries/lso_en.pdf). [Last accessed on 28 Feb 2017]
- World Health Organization. Non-communicable diseases progress monitor; 2015. [http://apps.who.int/iris/bitstream/10665/184688/1/9789241509459\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/184688/1/9789241509459_eng.pdf). [Last accessed on 28 Feb 2017].
- Renuga E, Ramakrishnan SR, Vanitha RN, Thenarasu P, Kannan G. Impact of continuous patient counselling on knowledge, attitude, and practices and medication adherence of diabetic patients attending outpatient pharmacy services. *Asian J Pharm Clin Res* 2016;9:364-9.
- Obirikorang Y, Obirikorang C, Anto EO, Acheampong E, Batu EN, Stella AD, et al. Knowledge of complications of diabetes mellitus among patients visiting the diabetes clinic at Sampa Government Hospital, Ghana: a descriptive study. *BMC Public Health* 2016;16:637.
- Adepu R, Madhu S. Influence of post-discharge counselling on health outcomes in diabetic and hypertensive patients. *Asian J Pharm Clin Res* 2011;4:28-33.
- Naveen B, Mahaboojan M, Padmanabha YR, Narayana G. Impact of clinical pharmacist mediated patient counselling on health-related quality of life in hypertensive patients. *Indian J Pharm Pract* 2014;7:34-40.
- Nnaemeka OR, Kingsley NC. Evaluation of patient's anti-diabetic medication counselling provided in a tertiary health care setting Nigeria. *Int Res J Pharm* 2012;3:319-23.
- Inamdar SZ, Kulkarni RV, Karajgi SR, Manvi FV, Ganachari MS, Mahendra Kumar BJ. Medication adherence in diabetes mellitus: an overview on pharmacist role. *Am J Adv Drug Delivery* 2013;1:238-50.
- Hughes JD, Wibowo Y, Sunderland B, Hoti K. The role of the pharmacist in the management of type 2 diabetes: current insights and future directions. *Int Pharm Res Pract J* 2017;6:15-27.
- Antoine SL, Pieper D, Mathes T, Eikermann M. Improving the adherence of type 2 diabetes mellitus patients with pharmacy care: a systematic review of the randomized rolled trial. *BMC Endocr Disord* 2014;14:53.
- Bollu M, Nalluri KK, Prakash AS, Lohith MN, Venkataramarao N. Study of knowledge, attitude, and practice of general population of Guntur toward silent killer diseases: hypertension and diabetes. *Asian J Pharm Clin Res* 2015;8:74-8.
- Akoko MB, Fon NP, Ngu CR, Ngu BK. Knowledge of hypertension and compliance with therapy among hypertensive patients in the Bamenda health district of Cameroon: a cross-sectional study. *Cardiol Ther* 2017;6:53-67.
- American Heart Association. Understanding blood pressure readings; 2016. [http://www.heart.org/HEARTORG/Conditions/HighBloodPressure/KnowYourNumbers/Understanding-Blood-Pressure-Readings\\_UCM\\_301764\\_Article.jsp#.Wdtj6l-CzIU](http://www.heart.org/HEARTORG/Conditions/HighBloodPressure/KnowYourNumbers/Understanding-Blood-Pressure-Readings_UCM_301764_Article.jsp#.Wdtj6l-CzIU). [Last accessed on 09 Oct 2017]
- Dekker MA, Amick EA, Scholcoff C, Doobay-Persaud A. A mixed-methods needs assessment of adult diabetes mellitus (type II) and hypertension care in Toledo, Belize. *BMC Health Serv Res* 2017;17:171.
- Nepal C, Vyas P, Bhattarai R, Acharya B, Thapa K, Shrestha SJ, et al. Knowledge, attitude and practice of type 2 diabetic patients of selected outreach clinic, Dharan, Nepal. *Al Ameen J Med Sci* 2017;10:78-85.
- The Global Diabetes Community. Blood sugar level ranges; 2017. [http://www.diabetes.co.uk/diabetes\\_care/blood-sugar-level-ranges.html](http://www.diabetes.co.uk/diabetes_care/blood-sugar-level-ranges.html). [Last accessed on 09 Oct 2017]
- Almas A, Godil SS, Lalani S, Samani AZ, Khan HA. Good knowledge about hypertension is linked to better control of hypertension; a multicentre cross-sectional study in Karachi, Pakistan. *BMC Res Notes* 2012;5:579.
- Familoni OB, Ogun AS, Aina OA. Knowledge and awareness of hypertension among patients with systemic hypertension. *J Natl Med Assoc* 2004;96:620-4.
- Fairchild CP, Nathan GA, Quinn M, Huang SE, Laiteerapong N. Patients' future expectations for diabetes and hypertension treatments: "through the Diet... I think this is going to go away. *J Gen Intern Med* 2016;32:49-55.
- Obirikorang Y, Obirikorang C, Anto OE, Acheampong E, Batu NE, Stella DA, et al. Knowledge of complications of diabetes mellitus among patients visiting the diabetes clinic at Sampa Government Hospital, Ghana: a descriptive study. *BMC Public Health* 2016;16:637.
- Adepu R, Somashekar MA. Influence of structured patient education on therapeutic outcomes in diabetes and hypertensive patients. *Asian J Pharm Clin Res* 2010;3:174-8.
- Abd El-Hay AS, El Mezayen ES. Knowledge and perceptions related to hypertension, lifestyle behaviour modifications and challenges that facing hypertensive patients. *J Nurs Health Sci* 2015;4:15-26.
- Bollampally M, Chandershekhar P, Kumar KP, Surakasula A, Srikanth S, Reddy TRM. Assessment of patient's knowledge, attitude and practice regarding hypertension. *Int J Res Med Sci* 2016;4:3299-304.
- Manobharathi M, Kalyani P, John William Felix A, Arulmani A. Factors associated with therapeutic non-compliance among type 2 diabetes mellitus patients in Chidambaram, Tamilnadu, India. *Int J Community Med Public Health* 2017;4:787-91.